



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

NATIONAL SENIOR CERTIFICATE

GRADE 12

MATHEMATICS P2

EXEMPLAR 2014

MEMORANDUM

MARKS: 150

This memorandum consists of 13 pages.

NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the marking memorandum.
- Assuming answers/values in order to solve a problem is NOT acceptable.

QUESTION 1

1.1	As the number of days that an athlete trained increased, the time taken to run the 100m event decreased. <p style="text-align: center;">OR</p> The fewer number of days an athlete trained, the longer the time he took to complete the 100m sprint. <p style="text-align: center;">OR</p> The greater number of days an athlete trained, the shorter the time he ran the 100m sprint.	✓ explanation (1)
1.2	(60 ; 18,1)	✓ (1)
1.3	$a = 17,81931464...$ $b = -0,070685358...$ $\therefore \hat{y} = -0,07x + 17,82$	✓✓ a ✓ b ✓ equation (4)
1.4	$\therefore \hat{y} \approx -0,07(45) + 17,82$ $\approx 14,67$ seconds	✓ substitution ✓ answer (2)
1.5	$r = -0,74$ (-0,740772594...)	✓✓ r (2)
1.6	There is a moderately strong relationship between the variables.	✓ moderately strong (1)
		[11]

QUESTION 2

<p>2.1</p>		<ul style="list-style-type: none"> ✓ grounding at 0 ✓ plotting at upper limits ✓ smooth shape of curve <p style="text-align: right;">(3)</p>
<p>2.2</p>	<p>$40 \leq t < 60$</p>	<ul style="list-style-type: none"> ✓ class <p style="text-align: right;">(1)</p>
<p>2.3</p>	<p>(96 ; 164) $\therefore 172 - 164 = 8$ learners</p>	<ul style="list-style-type: none"> ✓ 164 ✓ 8 <p style="text-align: right;">(2)</p>
<p>2.4</p>	<p>Frequency: 25; 44; 60; 28; 9; 6 $\text{Mean} = \frac{25 \times 10 + 44 \times 30 + 60 \times 50 + 28 \times 70 + 9 \times 90 + 6 \times 110}{172}$ $= \frac{8000}{172}$ $= 46,51 \text{ hours}$</p>	<ul style="list-style-type: none"> ✓ frequency ✓ midpoints ✓ $\frac{8000}{172}$ ✓ answer <p style="text-align: right;">(4) [10]</p>

QUESTION 3


3.1	$K(7 ; 0)$	✓ answer (1)
3.2	$1 = \frac{x_M + 7}{2}$ and $1 = \frac{y_M + 3}{2}$ $\therefore M(-5 ; -1)$	✓ x ✓ y (2)
3.3	$m_{PM} = \frac{3-1}{7-1}$ $= \frac{1}{3}$	✓ substitution ✓ answer (2)
3.4	$\tan \hat{P}SK = m_{PM} = \frac{1}{3}$ $\hat{P}SK = \tan^{-1}\left(\frac{1}{3}\right) = 18,43^\circ$ $\therefore \theta = 180^\circ - 90^\circ - 18,43^\circ = 71,57^\circ$	✓ $\tan \hat{P}SK = m_{PM}$ ✓ $\hat{P}SK$ ✓ θ (3)
3.5	$\cos 71,57^\circ = \frac{PK}{PS} = \frac{3}{PS}$ $PS = \frac{3}{\cos 71,57^\circ}$ $= 9,49$ units OR $\sin 18,43^\circ = \frac{PK}{PS} = \frac{3}{PS}$ $PS = \frac{3}{\sin 18,43^\circ}$ $= 9,49$ units	✓ correct ratio ✓ PS as subject ✓ answer (3) ✓ correct ratio ✓ PS as subject ✓ answer (3)
3.6	$N(x ; -2x + 17)$ $m_{TN} = m_{PM}$ (TN PM) $\frac{-2x + 17 - 5}{x - (-1)} = \frac{1}{3}$ $-6x + 36 = x + 1$ $-7x = -35$ $x = 5$ $\therefore y = -2(5) + 17 = 7$ $\therefore N(5 ; 7)$ OR	✓ N in terms of x ✓ equal gradients ✓ substitution ✓ x -value ✓ y -value (5)

	$m_{TM} = \frac{1}{3} \quad (\text{TN} \parallel \text{PM})$ <p>equation of TM:</p> $y - y_1 = \frac{1}{3}(x - x_1)$ $y - 5 = \frac{1}{3}(x - (-1))$ $y - 5 = \frac{1}{3}x + \frac{1}{3}$ $y = \frac{1}{3}x + 5\frac{1}{3}$ <p style="text-align: center;">OR</p> $y = \frac{1}{3}x + c$ $5 = \frac{1}{3}(-1) + c$ $5\frac{1}{3} = c$ $y = \frac{1}{3}x + 5\frac{1}{3}$ $-2x + 17 = \frac{1}{3}x + 5\frac{1}{3}$ $-2\frac{1}{3}x = -11\frac{2}{3}$ $x = 5$ $\therefore y = -2(5) + 17 = 7$ $\therefore N(5; 7)$	<p>✓ m_{TM}</p> <p>✓ equation of TM</p> <p>✓ equating</p> <p>✓ x-value</p> <p>✓ y-value</p> <p style="text-align: right;">(5)</p>
<p>3.7.1</p>	<p>$y = 5$</p>	<p>✓ equation</p> <p style="text-align: right;">(1)</p>
<p>3.7.2</p>	<p>gradient of AQ = $\tan 45^\circ$ or $\tan 135^\circ$ $= 1$ or -1</p> $m_{AQ} = \frac{5-1}{a-1} = \pm 1$ $\therefore a-1 = 4 \text{ or } -4$ $\therefore a = 5 \text{ or } -3$	<p>✓ $m_{AQ} = 1$ or</p> <p>✓ $m_{AQ} = -1$</p> <p>✓ substitution into gradient formula</p> <p>✓ x-value</p> <p>✓ y-value</p> <p style="text-align: right;">(5) [22]</p>

QUESTION 4

4.1	$M(-1 ; -1)$	✓ answer (1)
4.2	$m_{NT} = \frac{2-1}{3-4} = -1$ $\therefore m_{AT} = 1 \quad (\text{radius} \perp \text{tangent})$ $y - 1 = 1(x - 4)$ $y = x - 3$	✓ m_{NT} ✓ m_{AT} ✓ reason ✓ substitution of m and $(4 ; 1)$ ✓ equation (5)
4.3	$MR \perp AB$ (line from centre to midpt of chord) $MB^2 = MR^2 + RB^2$ (Theorem of Pythagoras) $9 = \left(\frac{\sqrt{10}}{2}\right)^2 + RB^2$ $RB^2 = \frac{13}{2}$ $RB = \sqrt{\frac{13}{2}}$ $AB = 2\left(\sqrt{\frac{13}{2}}\right) = \sqrt{26} \text{ units}$	✓ $MR \perp AB$ ✓ $MB = 3$ ✓ substitution into Theorem of Pythagoras ✓ AB in surd form (4)
4.4	$MN^2 = (-1 - 3)^2 + (-1 - 2)^2$ $= 16 + 9$ $= 25$ $MN = 5 \text{ units}$	✓ substitution into distance formula ✓ answer (2)
4.5	$r = 5 - 3 = 2 \text{ units}$ $\therefore (x - 3)^2 + (y - 2)^2 = 4$ $\therefore x^2 + y^2 - 6x - 4y + 9 = 0$	✓ r ✓ substitution into circle equation ✓ equation (3) [15]

QUESTION 5


5.1.1	$-\sin \alpha$ $= -\left(-\frac{4}{5}\right) = \frac{4}{5}$	✓ reduction ✓ answer (2)
5.1.2	$(-4)^2 + b^2 = 5^2$ $b^2 = 25 - 16 = 9$ $b = -3$ $\cos \alpha = \frac{-3}{5}$ 	✓ $b = -3$ ✓ answer (2)
5.1.3	$\sin(\alpha - 45^\circ)$ $= \sin \alpha \cos 45^\circ - \cos \alpha \sin 45^\circ$ $= -\frac{4}{5} \cdot \frac{1}{\sqrt{2}} - \left(-\frac{3}{5}\right) \cdot \frac{1}{\sqrt{2}}$ $= -\frac{1}{5\sqrt{2}}$ <p style="text-align: center;">OR</p> $\sin(\alpha - 45^\circ)$ $= \sin \alpha \cos 45^\circ - \cos \alpha \sin 45^\circ$ $= -\frac{4}{5} \cdot \frac{\sqrt{2}}{2} - \left(-\frac{3}{5}\right) \cdot \frac{\sqrt{2}}{2}$ $= -\frac{\sqrt{2}}{10}$	✓ expansion ✓ $\frac{1}{\sqrt{2}}$ ✓ answer in simplest form (3)
5.2.1	$LHS = \frac{8 \sin x \cdot \cos x}{\sin^2 x - \cos^2 x}$ $= \frac{4(2 \sin x \cdot \cos x)}{\sin^2 x - \cos^2 x}$ $= \frac{4 \sin 2x}{-(\cos^2 x - \sin^2 x)}$ $= \frac{4 \sin 2x}{-\cos 2x}$ $= -4 \tan 2x$	✓ $\sin x$ ✓ $\cos x$ ✓ $\cos^2 x$ ✓ $4 \sin 2x$ ✓ factorise ✓ $-\cos 2x$ (6)
5.2.2	Undefined when $\cos 2x = 0$ or $\tan 2x = \infty$: $x = 45^\circ$ and $x = 135^\circ$	✓ 45° ✓ 135° (2)

5.3	$1 - 2\sin^2 \theta + 4\sin^2 \theta - 5\sin \theta - 4 = 0$ $2\sin^2 \theta - 5\sin \theta - 3 = 0$ $(2\sin \theta + 1)(\sin \theta - 3) = 0$ $\therefore \sin \theta = -\frac{1}{2} \quad \text{or} \quad \sin \theta = 3 \text{ (no solution)}$ $\therefore \theta = 210^\circ + 360^\circ k \quad \text{or} \quad \theta = 330^\circ + 360^\circ k \quad ; k \in \mathbb{Z}$ <p>OR</p> $\therefore \theta = 210^\circ + 360^\circ k \quad \text{of} \quad \theta = 30^\circ + 360^\circ k \quad ; k \in \mathbb{Z}$	$\checkmark 1 - 2\sin^2 \theta$ \checkmark standard form \checkmark factors \checkmark no solution $\checkmark 210^\circ$ $\checkmark 330^\circ$ $\checkmark + 360^\circ k \quad ; k \in \mathbb{Z}$ (7) [22]
-----	--	--

QUESTION 6

6.1	$b = \frac{1}{2}$	\checkmark value of b (1)
6.2	A(30° ; 1)	$\checkmark 30^\circ$ $\checkmark 1$ (2)
6.3	$x = 160^\circ$	$\checkmark x = 160^\circ$ (1)
6.4	$h(x) = 2\cos(x - 30^\circ) + 1$ $y \in [-1 ; 3]$ OR $-1 \leq y \leq 3$	\checkmark critical values \checkmark notation (2) [6]

QUESTION 7

<p>7.1</p>	<p>Draw $CD \perp AB$ In $\triangle ACD$: $\sin A = \frac{CD}{b} \therefore CD = b \cdot \sin A$ In $\triangle CBD$: $\sin B = \frac{CD}{a} \therefore CD = a \cdot \sin B$ $\therefore b \cdot \sin A = a \cdot \sin B$ $\therefore \frac{\sin A}{a} = \frac{\sin B}{b}$</p> 	<p>✓ construction ✓ sin A ✓ making CD the subject ✓ sin B ✓ $b \cdot \sin A = a \cdot \sin B$ (5)</p>
<p>7.2.1</p>	<p>$\hat{S}PQ = 180^\circ - 2x$ (opp \angles of cyclic quad) $\hat{P}SQ + \hat{P}QS = 2x$ (sum of \angles in \triangle) $\hat{P}SQ = \hat{P}QS = x$ (\angles opp equal sides)</p>	<p>✓ $\hat{S}PQ = 180^\circ - 2x$ (S/R) ✓ reason (2)</p>
<p>7.2.2</p>	$\frac{\sin \hat{S}PQ}{\sin(180^\circ - 2x)} = \frac{\sin \hat{P}SQ}{\sin x}$ $\frac{SQ}{k} = \frac{PQ}{\sin x}$ $SQ = \frac{k \sin 2x}{\sin x}$ $SQ = \frac{k(2 \sin x \cdot \cos x)}{\sin x} = 2k \cos x$ <p style="text-align: center;">OR</p> $SQ^2 = PQ^2 + PS^2 - 2PQ \cdot PS \cdot \cos \hat{S}PQ$ $= k^2 + k^2 - 2 \cdot k \cdot k \cdot \cos(180^\circ - 2x)$ $= 2k^2 + 2k^2 \cos 2x$ $= 2k^2 + 2k^2(2\cos^2 x - 1)$ $= 4k^2 \cos^2 x$ $SQ = 2k \cos x$	<p>✓ substitution into correct formula ✓ sin 2x ✓ SQ subject ✓ $2 \sin x \cdot \cos x$ (4) ✓ substitution into correct formula ✓ $-\cos 2x$ ✓ $2\cos^2 x - 1$ ✓ simplification (4)</p>
<p>7.2.3</p>	$\tan y = \frac{3}{k}$ $k = \frac{3}{\tan y}$ $SQ = 2 \cos x \left(\frac{3}{\tan y} \right)$ $\therefore = \frac{6 \cos x}{\tan y}$	<p>✓ tan ratio ✓ k subject and substitution (2) [13]</p>

QUESTION 8


8.1	the angle subtended by the chord in the alternate segment	✓ correct theorem (1)
8.2.1	$\hat{B}_1 = \hat{E}_1 = 68^\circ$ (tan chord theorem)	✓ $\hat{E}_1 = 68^\circ$ ✓ reason (2)
8.2.2	$\hat{E}_1 = \hat{B}_3 = 68^\circ$ (alt \angle s; $AE \parallel BC$)	✓ $\hat{B}_3 = 68^\circ$ (S/R) (1)
8.2.3	$\hat{D}_1 = \hat{B}_3 = 68^\circ$ (ext \angle of cyclic quad)	✓ $\hat{D}_1 = 68^\circ$ ✓ reason (2)
8.2.4	$\hat{E}_2 = 20^\circ + 68^\circ$ $= 88^\circ$ (ext \angle of Δ)	✓ $\hat{E}_2 = 88^\circ$ (S/R) (1)
8.2.5	$\hat{C} = 180^\circ - 88^\circ$ $= 92^\circ$ (opp \angle s of cyclic quad)	✓ $\hat{C} = 92^\circ$ ✓ reason (2) [9]

QUESTION 9

<p>9.1</p>	<p>$\hat{D}_4 = \hat{A} = x$ (tan chord theorem) $\hat{A} = \hat{D}_2 = x$ (\angles opp equal sides)</p>	<p>✓ $\hat{A} = x$ ✓ reason ✓ $\hat{A} = \hat{D}_2 = x$ (S/R) (3)</p>
<p>9.2</p>	<p>$\hat{M}_1 = 2x$ (ext \angle of Δ) or (\angle at centre = $2\angle$ at circum) $\hat{M}\hat{D}E = 90^\circ$ (radius \perp tan) $\hat{M}_2 = 90^\circ - 2x$ $\therefore \hat{E} = 180^\circ - (90^\circ + 90^\circ - 2x)$ (sum of \angles in ΔMDE) $= 2x$ $\therefore CM$ is a tangent (converse tan chord theorem)</p>	<p>✓ $\hat{M}_1 = 2x$ (S/R) ✓ $\hat{M}\hat{D}E = 90^\circ$ (S/R) ✓ $\hat{E} = 2x$ ✓ reason (4)</p>
<p>9.3</p>	<p>$\hat{M}_3 = 90^\circ$ (EM \perp AC) $\hat{A}\hat{D}B = 90^\circ$ (\angle in semi-circle) $\therefore FMBD$ a cyclic quad (ext \angle of quad = int opp \angle) OR $\hat{E}\hat{M}C = 90^\circ$ (EM \perp AC) $\hat{A}\hat{D}B = 90^\circ$ (\angle in semi-circle) $\therefore FMBD$ a cyclic quad (opp \angles of quad supp)</p>	<p>✓ $\hat{M}_3 = 90^\circ$ ✓ $\hat{A}\hat{D}B = 90^\circ$ (S/R) ✓ reason (3) ✓ $\hat{E}\hat{M}C = 90^\circ$ ✓ $\hat{A}\hat{D}B = 90^\circ$ (S/R) ✓ reason (3)</p>
<p>9.4</p>	<p>$DC^2 = MC^2 - MD^2$ (Theorem of Pythagoras) $= (3BC)^2 - (2BC)^2$ (MB = MD = radii) $= 9BC^2 - 4BC^2$ $= 5BC^2$</p>	<p>✓ Th of Pythagoras ✓ substitution ✓ $9BC^2 - 4BC^2$ (3)</p>
<p>9.5</p>	<p>In ΔDBC and ΔDFM: $\hat{D}_4 = \hat{D}_2 = x$ (proven in 9.1) $\hat{B}_1 = \hat{F}_2$ (ext \angle of cyclic quad) $\hat{C} = \hat{M}_2$ $\therefore \Delta DBC \parallel \parallel \Delta DFM$ (\angle; \angle; \angle)</p>	<p>✓ $\hat{D}_4 = \hat{D}_2$ ✓ $\hat{B}_1 = \hat{F}_2$ ✓ reason ✓ $\hat{C} = \hat{M}_2$ or (\angle; \angle; \angle) (4)</p>
<p>9.6</p>	<p>$\frac{DM}{FM} = \frac{DC}{BC}$ ($\Delta DBC \parallel \parallel \Delta DFM$) $= \frac{\sqrt{5}BC}{BC}$ $= \sqrt{5}$</p>	<p>✓ S ✓ answer (2) [19]</p>

QUESTION 10

10.1



Construction: Join DC and BE and heights k and h

$$\frac{\text{area } \triangle ADE}{\text{area } \triangle DEB} = \frac{\frac{1}{2} \cdot AD \cdot k}{\frac{1}{2} \cdot DB \cdot k} = \frac{AD}{DB} \quad (\text{equal heights})$$

$$\frac{\text{area } \triangle ADE}{\text{area } \triangle DEC} = \frac{\frac{1}{2} \cdot AE \cdot h}{\frac{1}{2} \cdot EC \cdot h} = \frac{AE}{EC} \quad (\text{equal heights})$$

But Area $\triangle DEB = \text{Area } \triangle DEC$ (same base, same height)

$$\therefore \frac{\text{area } \triangle ADE}{\text{area } \triangle DEB} = \frac{\text{area } \triangle ADE}{\text{area } \triangle DEC}$$

$$\therefore \frac{AD}{DB} = \frac{AE}{EC}$$

✓ construction

$$\checkmark \frac{\text{area } \triangle ADE}{\text{area } \triangle DEB} = \frac{AD}{DB}$$

✓ reason

$$\checkmark \frac{\text{area } \triangle ADE}{\text{area } \triangle DEC} = \frac{AE}{EC}$$

✓ Area $\triangle DEB = \text{Area } \triangle DEC$ (S/R)

$$\checkmark \frac{\text{area } \triangle ADE}{\text{area } \triangle DEB} = \frac{\text{area } \triangle ADE}{\text{area } \triangle DEC}$$

(6)

<p>10.2.1</p>	$\frac{AB}{BE} = \frac{AC}{CD} \quad (\text{Prop Th; } BC \parallel ED)$ $\frac{1}{3} = \frac{3}{CD}$ $\therefore CD = 9 \text{ units}$	<p>✓ $\frac{AB}{BE} = \frac{AC}{CD}$ (S/R)</p> <p>✓ substitution</p> <p>✓ answer</p> <p>(3)</p>
<p>10.2.2</p>	$\frac{DG}{GA} = \frac{FD}{FE} \quad (\text{Prop Th; } FG \parallel EA)$ $\frac{9-x}{3+x} = \frac{3}{6}$ $54 - 6x = 9 + 3x$ $-9x = -45$ $x = 5$	<p>✓ $\frac{DG}{GA} = \frac{FD}{FE}$ (S/R)</p> <p>✓ substitution</p> <p>✓ simplification</p> <p>✓ answer</p> <p>(4)</p>
<p>10.2.3</p>	<p>In $\triangle ABC$ and $\triangle AED$:</p> <p>\hat{A} is common</p> <p>$\hat{ABC} = \hat{E}$ (corres \angles; $BC \parallel ED$)</p> <p>$\hat{ACB} = \hat{D}$ (corres \angles; $BC \parallel ED$)</p> <p>$\triangle ABC \sim \triangle AED$ (\angle, \angle, \angle)</p> <p>$\therefore \frac{BC}{ED} = \frac{AC}{AD}$</p> <p>$\frac{BC}{9} = \frac{3}{12}$</p> <p>$BC = 2\frac{1}{4} \text{ units}$</p>	<p>✓ \hat{A} is common</p> <p>✓ $\hat{ABC} = \hat{E}$ (S/R)</p> <p>✓ $\hat{ACB} = \hat{D}$ (S/R)</p> <p>or ($\angle; \angle; \angle$)</p> <p>✓ $\frac{BC}{ED} = \frac{AC}{AD}$</p> <p>✓ answer</p> <p>(5)</p>
<p>10.2.4</p>	$\frac{\text{area } \triangle ABC}{\text{area } \triangle GFD} = \frac{\frac{1}{2} AC \cdot BC \cdot \sin \hat{ACB}}{\frac{1}{2} GD \cdot FD \cdot \sin \hat{D}}$ $= \frac{\frac{1}{2} (3) (2\frac{1}{4}) \sin \hat{D}}{\frac{1}{2} (4) (3) \sin \hat{D}} \quad (\text{corres } \angle\text{s; } BC \parallel ED)$ $= \frac{9}{16}$	<p>✓ use of area rule</p> <p>✓ correct sides and angles</p> <p>✓ substitution of values</p> <p>✓ $\sin \hat{ACB} = \sin \hat{D}$ (S/R)</p> <p>✓ answer</p> <p>(5)</p> <p>[23]</p>

TOTAL: 150