

# basic education

Department: Basic Education **REPUBLIC OF SOUTH AFRICA** 

NATIONAL SENIOR CERTIFICATE

GRADE 12

## **MATHEMATICS P2**

\_ \_ \_ \_ \_ \_ \_ \_ \_ \_ \_ \_ \_ \_ \_ \_

# FEBRUARY/MARCH 2015

............

**MARKS: 150** 

TIME: 3 hours

This question paper consists of 14 pages, 5 diagram sheets and 1 information sheet.



#### **INSTRUCTIONS AND INFORMATION**

Read the following instructions carefully before answering the questions.

- 1. This question paper consists of 10 questions.
- 2. Answer ALL the questions.
- 3. Clearly show ALL calculations, diagrams, graphs, et cetera which you have used in determining your answers.
- 4. Answers only will NOT necessarily be awarded full marks.
- 5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
- 6. If necessary, round off answers to TWO decimal places, unless stated otherwise.
- 7. FIVE diagram sheets for QUESTIONS 1.3, 7, 8, 9.2, 9.3 and 10 are attached at the end of this question paper. Write your centre number and examination number on these sheets in the spaces provided and insert them inside the back cover of your ANSWER BOOK.
- 8. Diagrams are NOT necessarily drawn to scale.
- 9. Number the answers correctly according to the numbering system used in this question paper.
- 10. Write neatly and legibly.



## **QUESTION 1**

The table below shows the distances (in kilometres) travelled daily by a sales representative for 21 working days in a certain month.

131	132	140	140	141	144	146
147	149	150	151	159	167	169
169	172	174	175	178	187	189

1.1	Calculate t	he mean distance travelled by the sales representative.	(2)
1.2	Write dow	n the five-number summary for this set of data.	(4)
1.3	Use the sc this set of	aled line on DIAGRAM SHEET 1 to draw a box-and-whisker diagram for data.	(2)
1.4	Comment	on the skewness of the data.	(1)
1.5	Calculate 1	the standard deviation of the distance travelled.	(2)
1.6	The sales on each of in terms of	representative discovered that his odometer was faulty. The actual reading $f$ the 21 days was $p$ km more than that which was indicated. Write down, f $p$ (if applicable), the:	
	1.6.1	Actual mean	(1)
	1.6.2	Actual standard deviation	(1) [ <b>13</b> ]



An ice-cream shop recorded the sales of ice cream, in rand, and the maximum temperature, in °C, for 12 days in a certain month. The data that they collected is represented in the table and scatter plot below.

Temperature in °C	24,2	26,4	21,9	25,2	28,5	32,1	29,4	35,1	33,4	28,1	32,6	27,2
Sales of ice cream in rand	215	325	185	332	406	522	412	614	544	421	445	408



2.1	Describe the influence of temperature on the sales of ice cream in the scatter plot.	(1)
2.2	Give a reason why this trend cannot continue indefinitely.	(1)
2.3	Calculate an equation for the least squares regression line (line of best fit).	(4)
2.4	Calculate the correlation coefficient.	(1)
2.5	Comment on the strength of the relationship between the variables.	(1) [8]



In the diagram below points P(5; 13), Q(-1; 5) and S(7,5; 8) are given. SR || PQ where R is the *y*-intercept of SR. The *x*-intercept of SR is B. QR is joined.





4.1 In the diagram below, the circle centred at M(2; 4) passes through C(-1; 2) and cuts the y-axis at E. The diameter CMD is drawn and ACB is a tangent to the circle.



$(x-5)^2 +$	$(y+1)^2 = 9$ will intersect. Show ALL calculations.	(6) [ <b>22</b> ]
Determine	whether or not the circles having equations $(x+2)^2 + (y-4)^2 = 25$ and	
4.1.5	Show that EM is parallel to AB.	(2)
4.1.4	Calculate the coordinates of E.	(4)
4.1.3	Determine the equation of AB in the form $y = mx + c$ .	(5)
4.1.2	Write down the coordinates of D.	(2)
4.1.1	Determine the equation of the circle in the form $(x-a)^2 + (y-b)^2 = r^2$ .	(3)



4.2

(6)

#### 7 NSC

#### **QUESTION 5**

- 5.1 If  $x = 3 \sin \theta$  and  $y = 3 \cos \theta$ , determine the value of  $x^2 + y^2$ . (3)
- 5.2 Simplify to a single term:

$$\sin(540^\circ - x) \cdot \sin(-x) - \cos(180^\circ - x) \cdot \sin(90^\circ + x)$$

5.3 In the diagram below, T(x; p) is a point in the third quadrant and it is given that  $\sin \alpha = \frac{p}{\sqrt{1+p^2}}$ .



5.3.1 Show that 
$$x = -1$$
. (3)

5.3.2 Write  $\cos(180^\circ + \alpha)$  in terms of p in its simplest form. (2)

5.3.3 Show that  $\cos 2\alpha$  can be written as  $\frac{1-p^2}{1+p^2}$ . (3)

5.4 5.4.1 For which value(s) of x will  $\frac{2\tan x - \sin 2x}{2\sin^2 x}$  be undefined in the interval  $0^\circ \le x \le 180^\circ$ ? (3)

5.4.2 Prove the identity: 
$$\frac{2\tan x - \sin 2x}{2\sin^2 x} = \tan x$$
(6)  
[26]



6.1 In the figure, points K, A and F lie in the same horizontal plane and TA represents a vertical tower.  $\hat{ATK} = x$ ,  $\hat{KAF} = 90^\circ + x$  and  $\hat{KFA} = 2x$  where  $0^\circ < x < 30^\circ$ . TK = 2 units.



6.1.1	Express AK in terms of $\sin x$ .		(2)
6.1.2	Calculate the numerical value of KF.		(5)



6.2 In the diagram below, a circle with centre O passes through A, B and C. BC = AC = 15 units. BO and OC are joined. OB = 10 units and BOC = x.



#### Calculate:

6.2.1	The size of $x$	(4)
6.2.2	The size of ACB	(3)
6.2.3	The area of $\triangle ABC$	(2) [ <b>16</b> ]





## GIVE REASONS FOR YOUR ANSWERS IN QUESTIONS 7, 8, 9 AND 10.

#### **QUESTION 7**

In the diagram, AB is a chord of the circle with centre O. M is the midpoint of AB. MO is produced to P, where P is a point on the circle. OM = x units, AB = 20 units and  $\frac{PM}{OM} = \frac{5}{2}$ .



7.1	Write down the length of MB.	(1)
7.2	Give a reason why $OM \perp AB$ .	(1)
7.3	Show that $OP = \frac{3x}{2}$ units.	(2)
7.4	Calculate the value of $x$ .	(3)

Copyright reserved



[7]

In the diagram below, the circle with centre O passes through A, B, C and D.

AB || DC and  $\hat{BOC} = 110^{\circ}$ .

The chords AC and BD intersect at E.

EO, BO, CO and BC are joined.



8.1

Calculate the size of the following angles, giving reasons for your answers:

8.1.1 D	(2)
8.1.2 Â	(2)
8.1.3 Ê <sub>2</sub>	(4)

8.2 Prove that BEOC is a cyclic quadrilateral. (2) [10]



(1)

12 NSC

#### **QUESTION 9**

9.1 Complete the statement of the following theorem:

The exterior angle of a cyclic quadrilateral is equal to ...

9.2 In the diagram below the circle with centre O passes through points S, T and V. PR is a tangent to the circle at T. VS, ST and VT are joined.



Given below is the partially completed proof of the theorem that states that  $V\hat{T}R = \hat{S}$ . Using the above diagram, complete the proof of the theorem on DIAGRAM SHEET 3.

Construction: Draw diameter TC and join CV.

Statement	Reason
<b>Let:</b> $V\hat{T}R = \hat{T}_1 = x$	7
$\hat{\mathbf{V}}_1 + \hat{\mathbf{V}}_2 = \dots$	
$\hat{\mathrm{T}}_2 = 90^{\circ} - x$	
∴ Ĉ =	Sum of the angles of a triangle
$\therefore \hat{\mathbf{S}} = x$	
$\therefore \dot{V}\hat{T}R = \hat{S}$	

(5)

Copyright reserved

9.3 In the figure, TRSW is a cyclic quadrilateral with TW = WS. RT and RS are produced to meet tangent VWZ at V and Z respectively. PRQ is a tangent to the circle at R. RW is joined.  $\hat{R}_2 = 30^\circ$  and  $\hat{R}_4 = 50^\circ$ .



9.3.4	Prove that $WR^2 = RV \times RS$ .	(5) [ <b>22</b> ]
	(b) Ŷ	(4)
	(a) $\hat{S}_2$	(3)
9.3.3	Determine, with reasons, the size of:	
9.3.2	State, with reasons, TWO other angles equal to 30°.	(3)
9.3.1	Give a reason why $R_3 = 30^{\circ}$ .	(1)



In  $\Delta$ TRM,  $\hat{M} = 90^{\circ}$ . NP is drawn parallel to TR with N on TM and P on RM. It is further given that RT = 3PN.



10.1 Give reasons for the statements below. Use DIAGRAM SHEET 5.

	Statement	Reason	
-	In $\Delta$ PNM and $\Delta$ RTM		
10.1.1	$\hat{N}_1 = \hat{T}$		
	is common		
10.1.2	$\therefore \Delta PNM     \Delta RTM$		(2)

10.2 Prove that  $\frac{PM}{RM} = \frac{1}{3}$ . (2)

10.3 Show that  $RN^2 - PN^2 = 2RP^2$ .

(4) [8]

**TOTAL: 150** 





## **QUESTION 7**





Mathematics/P2

DBE/Feb.-Mar. 2015





Mathematics/P2



NSC

**DIAGRAM SHEET 3** 

**QUESTION 9.2** 



Construction: Draw diameter CT and join CV.

Statement	Reason
Let: $\hat{VTR} = \hat{T}_1 = x$	
$\hat{\mathbf{V}}_1 + \hat{\mathbf{V}}_2 = \dots$	е 
$\hat{\mathrm{T}}_2 = 90^{\circ} - x$	· · · · · · · · · · · · · · · · · · ·
$\therefore \hat{C} = \dots$	Sum of the angles of a triangle
$\therefore \hat{S} = x$	
$\therefore \dot{V}\hat{T}R = \hat{S}$	



Mathematics/P2

DBE/Feb.-Mar. 2015

Q



3

2

 $\overline{2}/S$ 

NSC



4

<b>CENTRE NUMBER:</b>						5				
EXAMINATION NUMBER:						-				
DIAGRAM SHEET 5										
<b>QUESTION 10</b>						-	Г			
					/					
				/						
			/							
		*					N			
	/				/	/1				
				/	1					

R

NSC

10.1

	Statement	Reason				
	In $\triangle PNM$ and $\triangle RTM$ :					
10.1.1	$\hat{N}_1 = \hat{T}$					
	$\hat{M}$ is common					
10.1.2	$\therefore \Delta PNM     \Delta RTM$					

EASTERN CAPE

P

M

#### **INFORMATION SHEET**

$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$				
A = P(1+ni)	A = P(1 - ni)		$A = P(1-i)^n$	$A = P(1+i)^n$
$T_n = a + (n-1)d$		$\mathbf{S}_n = \frac{1}{2}$	$\frac{n}{2}[2a+(n-1)d]$	
$T_n = ar^{n-1}$		$S_n = -$	$\frac{a(r^n-1)}{r-1}  ; r \neq 1$	$S_{\infty} = \frac{a}{1-r}; -1 < r < 1$
$F = \frac{x\left[(1+i)^n - 1\right]}{i}$		$P = \frac{x}{2}$	$\frac{\left[1-\left(1+i\right)^{-n}\right]}{i}$	
$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x+h) - f(x+h) - f(x+h)}{h}$	f(x)			
$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_2)^2}$	$\overline{(v_1)^2}$	$M\left(\frac{x_1}{x_1}\right)$	$\frac{+x_2}{2}; \frac{y_1+y_2}{2}$	
y = mx + c	$y - y_1 = m(x - x)$	$-x_{1})$	$m = \frac{y_2 - y_1}{x_2 - x_1}$	$m = \tan \theta$
$(x-a)^2 + (y-b)^2 = r^2$				
In $\triangle ABC$ : $\frac{a}{\sin A} = \frac{b}{\sin B}$	$=\frac{c}{\sin C}$			
$a^2 = b^2 + c^2$	-2 <i>bc</i> .cos <i>A</i>			
area $\Delta ABC$ =	$=\frac{1}{2}ab.\sin C$			
$\sin(\alpha+\beta)=\sin\alpha.\cos\beta$	$+\cos\alpha.\sin\beta$		$\sin(\alpha-\beta)=\sin\alpha.co$	osβ–cosα.sinβ
$\cos(\alpha+\beta)=\cos\alpha.\cos\beta$	$\beta - \sin \alpha . \sin \beta$		$\cos(\alpha-\beta)=\cos\alpha.$	$\cos\beta + \sin\alpha . \sin\beta$
$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$	χ		$\sin 2\alpha = 2\sin \alpha .\cos \beta$	α
$\overline{x} = \frac{\sum fx}{n}$			$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \overline{x})^2}{n}$	
$P(A) = \frac{n(A)}{n(S)}$			P(A  or  B) = P(A) +	P(B) - P(A  and  B)

 $b = \frac{\sum (x - \overline{x})(y - \overline{y})}{\sum (x - \overline{x})^2}$ 

Copyright reserved

 $\hat{y} = a + bx$ 

